

Restylane[®]


SILK

DO YOU WANT FULLER, YOUNGER-LOOKING LIPS BUT FEAR THE FAKE LOOK?

Fear no more, and say hello to natural-looking lips with *Restylane[®] Silk*—the one and only FDA-approved lip and line filler that delivers the smooth and subtle look you want.

As we get older, the lines around the mouth show signs of aging. This can result in lip thinning, lost shape, and an increase in vertical lines above the lip that cause lipstick to feather. *Restylane Silk* provides natural-looking results in these areas by using a refined, smooth gel formulation.

See the difference *Restylane Silk* made in this patient



Before

After

EXPERIENCE THE DIFFERENCE FOR YOURSELF

Contact our office today to schedule your *Restylane Silk* consultation and become one step closer to achieving the lips you desire.

To learn more about *Restylane Silk*, visit RestylaneSilk.com

Please see the *Instructions for Use* at RestylaneSilk.com or call 1-855-425-8722. For more information, talk to your doctor.

Important Safety Information

Indication: *Restylane[®] Silk* is indicated for submucosal implantation for lip augmentation and dermal implantation for correction of perioral rhytids in patients over the age of 21.

Restylane Silk contains traces of gram-positive bacterial protein and is contraindicated for patients with allergies to such material or for patients with severe allergies that have required in-hospital treatment. *Restylane Silk* should not be used by patients with bleeding disorders, with hypersensitivity to amide-type local anesthetics, such as lidocaine, under the age of 22, or by women who are pregnant or breastfeeding. *Restylane Silk* should not be injected anywhere except the dermis or lip submucosa.

Use of *Restylane Silk* at the site of skin sores, pimples, rashes, hives, cysts, or infection should be postponed until healing is complete. The most commonly observed side effects are swelling, tenderness, bruising, pain, and redness at the injection site. These are typically mild in severity and resolve in 2-7 days after treatment. Serious but rare side effects include delayed onset infections, recurrence of herpetic eruptions, and superficial necrosis and scarring at the injection site. Do not implant into blood vessels. Use with caution in patients recently treated with anticoagulant or platelet inhibitors to avoid bleeding and bruising.

Treatment volume should be limited to 1.5 mL per lip per treatment and 1.0 mL for perioral rhytid correction, as greater amounts significantly increase moderate and severe injection site reactions. The safety or effectiveness of treatment in areas other than lips and perioral rhytids has not been established in controlled clinical studies.

Restylane Silk is only available through a licensed practitioner. Complete *Instructions for Use* are available at www.RestylaneSilk.com.



All trademarks are the property of their respective owners.
© 2015 Galderma Laboratories, L.P.
SLK-156A 06/15